

(19)



Europäisches Patentamt

European Patent Office

Office européen des brevets



(11)

EP 0 921 192 A1

(12)

## EUROPEAN PATENT APPLICATION

(43) Date of publication:

09.06.1999 Bulletin 1999/23

(51) Int. Cl.<sup>6</sup>: C12N 15/12, C07K 14/47,

A61K 38/17, C12Q 1/68,

C12N 1/19, C12N 1/21

(21) Application number: 97203781.6

(22) Date of filing: 03.12.1997

(84) Designated Contracting States:

AT BE CH DE DK ES FI FR GB GR IE IT LI LU MC  
NL PT SE

Designated Extension States:

AL LT LV MK RO SI

(71) Applicant: Leadd B.V.

2333 AL Leiden (NL)

(72) Inventors:

- Noteborn, Mathieu Hubertus Maria  
2352 EH Leiderdorp (NL)

- Danen - van Oorschot, Astrid A. A. M.  
2651 VG Berkel en Rodenrijs (NL)

(74) Representative:

Smulders, Theodorus A.H.J., Ir.  
Vereenigde Octroolbureaux  
Nieuwe Parklaan 97  
2587 BN 's-Gravenhage (NL)

Remarks:

The applicant has subsequently filed a sequence  
listing and declared, that it includes no new matter.

## (54) Molecules Interacting with apoptin

(57) The invention relates to activation of apoptosis  
by means of interference of Hou-like and/or IFP35-like  
compounds.

Also the invention relates to anti-tumor therapies  
with compounds, which negatively interfere with Hou-  
like and/or IFP35-like compounds leading to induction  
of apoptosis, resulting in the elimination of tumor cells.

Also the invention relates to therapies for diseases  
related to aberrant apoptosis induction, such as auto-  
immune disease.

Also the invention describes the diagnosis of cells,  
which are susceptible to apoptin- or apoptin-like  
induced apoptosis.

new treatments and diagnosis for diseases related with aberrancies in the apoptotic process, such as cancer and autoimmune diseases.

[0013] Proteins found associating with apoptin include members of the family of Nmi/Hou-like and IFP-like proteins.

[0014] Thus the invention provides a recombinant and/or isolated nucleic acid molecule encoding at least a functional part of a member of the family of Nmi-like proteins or at least a functional part of a member of the family of Hou-like proteins or at least a functional part of a member of the family of IFP35-like proteins for use in the induction of apoptosis in a population of cells related to a pathological condition.

[0015] As explained herein the expression of Hou is connected to oncogenes and has been found to be high in certain transformed cells. These are typically the cells that can be induced to go into apoptosis by apoptotic agents such as apoptin. Typically providing a cell with Hou-like activity will therefor increase the chance of inducing apoptosis in such a cell. IFP35-like proteins are involved in transporting apoptotic substances to the nucleus of cells. Under influence of for instance interferons these proteins localize in the nucleus. Therefor IFP-like activity is used to get apoptin-like activity into the nucleus, which is important for the induction of apoptosis, for instance through Hou-like proteins. The Hou-like activity or Nmi-like activity is defined herein as any molecule capable of exerting the same or a similar function as the original Hou-like (Nmi-like) protein. The same definition goes for IFP-activity. Typically such a molecule can be encoded by a nucleic acid molecule which comprises at least a functional and specific part of the sequence of figure 1, 2, 4 or 5 or encoding an amino sequence of figure 6 or a sequence at least 60, preferably 70, preferably 90 % homologous with said functional and specific sequence or comprising a sequence hybridizing to any of the foregoing sequences under stringent conditions. In order to be able to express the Hou-like activity and/or the IFP-like activity it is preferred to have an expression vector encoding said activity. Expression vectors are nucleic acid molecules which can be brought into cells, or transfect cells themselves and which have the machinery (together with the machinery of the host cell) to express proteins encoded on the expression vector when present in a cell.

[0016] It is preferred that cells which are provided, according to the invention, with Hou-like activity and/or IFP-like activity, are also provided with apoptosis inducing activity, preferably apoptin-like activity, which is defined along the same lines as Hou-like activity. In order to get the activity into the cells in which apoptosis has to be induced it is possible and preferred to use a gene delivery vehicle. A gene delivery vehicle is a means to transport a nucleic acid molecule capable of expressing the wanted activity in a host cell into said host cell. Gene delivery vehicles are known in the art. They include for instance recombinant viruses such as adenoviruses and retroviruses, but also non-viral vehicles such as polymers and liposomes have been suggested. Methods of targeting gene delivery vehicles to target cells are also known in the art and need not be elaborated herein. The invention also provides the newly identified molecules themselves, both the nucleic acid molecules (meaning DNA coding and/or non coding strands as well as RNA) and the proteinaceous molecules (peptides, polypeptides, glycoproteins and associations between proteins and RNA's and the like). Based on the given sequences other familymembers of the Hou/Nmi and IFP families will be identified having the same or similar function. Typically such molecules will have high homology to the sequences given herein.

[0017] For nucleic acid molecules the homology is expected to be at least 60, preferably 70, more preferably 80% therewith.

[0018] These nucleic acid molecules can of course again be incorporated into expression vectors as mentioned hereinbefore. Preferably these expression vectors also encode apoptotic activity, preferably apoptin or a functional fragment and/or equivalent thereof.

[0019] These expression vectors can again be made into gene delivery vehicles.

[0020] The invention also provides the recombinant or isolated proteinaceous substance comprising at least a functional part of a member of the family of Nmi/Hou-like proteins or at least a functional part of a member of the family of Hou-like proteins for use in the induction of apoptosis in a population of cells related to a pathological condition and an Nmi/Hou-like proteinaceous substance having at least a functional and/or specific part of the sequence of figure 3 or being encoded by a functional and/or specific part of the sequence of figure 1 or figure 2 or being at least 60, preferably 70, preferably 80% homologous to at least a functional and/or specific part of the sequence of figure 3 or being at least 60, preferably 70, preferably 80% homologous to a protein encoded by at least a functional and/or specific part of the sequence of figure 1 or figure 2 and an IFP35-like proteinaceous substance having at least a functional and/or specific part of the sequence of figure 6 or 7 or being encoded by a functional and/or specific part of the sequence of figure 4 or figure 5 or being at least 60, preferably 70, preferably 80% homologous to at least a functional and/or specific part of the sequence of figure 6 or 7 or being at least 60, preferably 70, preferably 80% homologous to a protein encoded by at least a functional and/or specific part of the sequence of figure 4 or figure 5.

[0021] A functional part in this document means having the same or similar activity (although the amount of activity may differ) A specific part herein means a part of sufficient size to be specific for the protein or nucleic acid or to be of sufficient size to distinguish the protein from another protein immunologically. The proteins disclosed herein can for instance also be used to identify further components of the apoptotic pathway.

[0022] The reason for bringing IFP-like activity and/or Hou-like activity together with apoptotic activity is of course to induce aberrant cells to go into apoptosis. Thus the invention also provides a method for inducing apoptosis in cells

**GAL4-activation domain-tagged cDNA library**

[0043] The expression vector pACT, containing the cDNAs from Epstein-Barr-virus-transformed human B cells fused to the GAL4 transcriptional activation domain, was used for detecting apoptin-associating proteins. The pACT c-DNA library is derived from the lambda-ACT cDNA library, as described by Durfee et al. 1993.

**Bacterial and Yeast strains**

[0044] The E.coli strain JM109 was the transformation recipient for the plasmid pGBT9 and pGBT-VP3. The bacterial strain electromax/DH10B was used for the transformation needed for the recovery the apoptin-associating pACT-cDNAs, and was obtained from GIBCO-BRL, USA.

[0045] The yeast strain Y190 was used for screening the cDNA library, and all other transformations which are part of the used yeast-two-hybrid system.

**Media**

[0046] For drug selections Luria Broth (LB) plates for E.coli were supplemented with ampicillin (50 microgram per ml). Yeast YPD and SC media were prepared as described by Rose et al. (1990).

**Transformation of competent yeast strain Y190 with plasmids pGBT-VP3 and pACT-cDNA and screening for beta-galactosidase activity.**

[0047] The yeast strain Y190 was made competent and transformed according to the methods described by Klebe et al. (Klebe et al., 1983). The yeast cells were first transformed with pGBT-VP3 and subsequently transformed with pACT-cDNA, and these transformed yeast cells were grown on histidine-minus plates, also lacking leucine and tryptophan.

[0048] Hybond-N filters were layed on yeast colonies, which were histidine-positive and allowed to wet completely. The filters were lifted and submerged in liquid nitrogen to permeabilize the yeast cells. The filters were thawed and layed with the colony side up on Whatman 3MM paper in a petridish with Z-buffer (Per liter: 16.1 gr  $\text{Na}_2\text{HPO}_4 \cdot 7\text{H}_2\text{O}$ , 5.5 gr  $\text{NaH}_2\text{PO}_4 \cdot \text{H}_2\text{O}$ , 0.75 gr KCl and 0.246 gr  $\text{MgSO}_4 \cdot 7\text{H}_2\text{O}$ , pH 7.0) containing 0.27% beta-mercapto-ethanol and 1 mg/ml X-gal. The filters were incubated for at least 15 minutes or during night.

**Recovery of plasmids from yeast**

[0049] Total DNA from yeast cells, which were histidine- and beta-galactosidase-positive, was prepared by using the glucolase-alkaline lysis method as described by Hoffman and Winston (1987) and used to transform Electromax/DH10B bacteria via electroporation using a Bio-Rad GenePulser according the manufacturer's specifications.

[0050] Transformants were plated on LB media containing ampicillin.

**Isolation of apoptin-associating pACT clones**

[0051] By means of colony-filter assay the colonies were lysed and hybridized to a radioactive-labeled 17-mer oligomer, which is specific for pACT (see also section Sequence analysis).

[0052] Plasmid DNA was isolated from the pACT-clones, and by means of XhoI digestion analysed for the presence of a cDNA insert.

**Sequence analysis**

[0053] The subclones containing the sequences encoding apoptin-associating proteins were sequenced using dideoxy NTPs according to the Sanger method which was performed by Eurogentec, Nederland BV (Maastricht, The Netherlands). The used sequencing primer was a CT-specific 17-mer comprising of the DNA-sequence 5'-TACCACTACAATGGATG-3'.

[0054] The sequences of the apoptin-associating proteins were compared with known gene sequences from the EMBL/Genbank.

**Results and discussion**

[0055] Apoptin induces specifically apoptosis in transformed cells, such as cell lines derived from human tumors. To identify the essential compounds in this cell-transformation-specific and/or tumor-specific apoptosis pathway, a yeast

Nmi, or Hou will be interchangeably used.

[0070] In this respect, the pattern of Nmi expression is interesting, since it is expressed at low levels in normal tissues, in contrast to its high levels of expression in transformed cell lines. Among eight cancer lines tested, highest levels were observed in four leukemia cell lines (Bao and Zervos, 1996).

[0071] In leukemias, a high expression of C-myc correlates with a high level of Nmi (HL-60, K562 and MOLT-4). The Nmi gene is located on chromosome 22, which is also involved in the t (9;22) translocation leading to the Bcr-Abl fusion protein, as seen in some leukemias (Rabbits, 1991, Sawyers and Dery, 1994).

[0072] Using a yeast genetic screen, Nmi was identified as a protein that binds to N-myc and C-myc. Myc proteins are important in the regulation of cell proliferation and differentiation. Together with ras or raf, myc can transform primary cells in culture. Nmi/Hou-like proteins will up-regulate the activity of Myc proteins via binding to them.

[0073] Up-regulation of Myc proteins has been described for Burkitt lymphomas, neuroblastomas and small cell lung carcinomas. Myc proteins contain a basic region, a helix-loop-helix (HLH) and a leucine zipper (Zip), and form homo- or heterodimers that can bind to specific DNA sequences and regulate transcription. Myc also forms heterodimers with Max. Myc/Max heterodimers activate transcription, whereas Max homodimers repress transcription, thus antagonizing Myc's function (Evan and Littlewood, 1993).

[0074] Nmi was found to interact with N-myc, c-myc, Max, Mxi1 and other transcription factors that have HLH and/or Zip motifs. Interaction with N-myc and C-myc was confirmed by co-precipitation experiments (Bao and Zervos, 1996).

#### Induction of apoptosis through interference with the function of Nmi/Hou-like proteins.

[0075] Our results indicate that apoptin can change the Nmi/Hou-like-mediated proliferation (transformation/tumor-formation) activity into a Nmi/Hou-like-mediated apoptotic activity. Remarkably, this Nmi/Hou-like-mediated apoptotic activity will be specific for transformed/tumor cells, due to the very high level of Nmi/Hou in transformed cells in combination with over-expression of (proto-)oncogenes, such as Myc.

[0076] By means of transient transfection assays, it was shown that over-expression of the determined Hou-like protein (see Fig. 3) and apoptin did result in induction of apoptosis in normal VH10-, VH25-fibroblasts. In contrast to normal fibroblasts which over-expressed only apoptin. This result indicates that Hou-like proteins are an important factor in (apoptin-induced) apoptosis.

[0077] The presented data imply that interference with the function of Nmi/Hou-like proteins resulting in apoptosis can be used as a specific anti-tumor therapy, or therapies of related diseases, such as auto-immune diseases.

#### Characteristics of the apoptin-associating protein IFP35

[0078] The other apoptin-associating protein is IFP35, which is an interferon(IFN)-induced leucine zipper protein of 282 a.a., and has an apparent molecular mass of 35 kD. It was isolated by differential screening from HeLa cells that had been treated with IFN- $\gamma$  (Bange et al., 1994).

[0079] IFP35 mRNA could be induced by IFN- $\gamma$  in different human cell types, including fibroblasts, macrophages, and epithelial cells. It has a leucine zipper motif at the N-terminus, but it lacks an adjacent basic domain required for DNA binding. It has been suggested that these types of proteins negatively regulate bZIP transcription factors by forming non-functional heterodimers. IFP35 was shown to form homodimers (Bange et al., 1994).

#### Induction of apoptosis by interference of IFP35 in combination with Hou/Nmi-like proteins.

[0080] IFP35 is found in the cell nucleus, after interferon treatment and is expressed in a wide variety of cell types including fibroblasts, macrophages and epithelial cells (Bange et al., 1994).

[0081] In general, virus infections trigger interferon production. It is likely that a CAV infection and/or expression of apoptin will result in interferon up-regulation, which might result in the translocation of IFP35 or IFP35-like proteins into the nucleus. IFP35 will transport apoptin also to the nucleus, due to its association.

[0082] It seems likely that if apoptin is transported into the nucleus by IFP35 it will be able to associate with the IFP35-homologous region within Hou/Nmi-like proteins. This association will cause an aberrant regulation of Hou/Nmi-regulated genes, such as the oncogene Myc. Subsequently, the cells over-expressing Nmi/Hou-like proteins and oncogenes, such as Myc will undergo apoptosis.

[0083] Experimental evidence for IFP35 as an essential factor in (apoptin) apoptosis induction was derived from the following experiments. Normal VH10 cells over-expressing Hou/Nmi, IFP35 and apoptin underwent faster apoptosis than normal VH10 cells expressing Hou/Nmi and apoptin.

retinoblastoma protein associates with the protein phosphate type I catalytic subunit. *Genes and Development* 7, 555-569.

8. Earnshaw, W.C., 1995. Nuclear changes in apoptosis. *Current Opinion in Cell Biology* 7, 337-343.

9. Evan, G.I. and Littlewood, T.D. (1993). *Current Opinions in Genetics Development* 3, 44-49.

10. Fields, S. and Song, O.K. (1989). A novel genetic system to detect protein-protein interactions. *Nature* 340, 245-246.

11. Hockenberry, D.M. (1994). Bcl-2 in cancer, development and apoptosis. *Journal of Cell Science, Supplement* 18, 51-55.

12. Hoffman, C.S. and Winston, F. (1987). A ten-minute DNA preparation from yeast efficiently releases autonomous plasmids for transformation of *Escherichia coli*. *Gene* 57, 267-272.

13. Kerr, J.F.R., Winterford, C.M., and Harmon, B.V. (1994). Apoptosis: Its significance in cancer and cancer therapy. *Cancer* 73, 2013-2026.

14. Klebe, R.J., Hariss, J.V., Sharp, Z.D., and Douglas, M.G. (1983). A general method for polyethylene-glycol-induced genetic transformation of bacteria and yeast. *Gene* 25, 333-341.

15. Levine, A.J. (1997). p53, the cellular gatekeeper for growth and division. *Cell* 88, 323-331.

16. Maniatis, T., Fritsch, E.F., and Sambrook, J. (1982). *Molecular Cloning: A Laboratory Manual*. CSHL Press, New York, USA.

17. Matsudaira, P. (1994). Actin crosslinking proteins at the leading edge. *Seminars in Cell biology* 5, 165-174.

18. McDonnell T.J., Meyn, R.E., Robertson, L.E. (1995). Implications of apoptotic cell death regulation in cancer therapy. *Seminars in Cancer Biology* 6, 53-60.

19. Noteborn, M.H.M. (1996). PCT application WO 96/41191. Apoptin induces apoptosis in human transformed and malignant cells but not in normal cells as essential characteristic for the development of an anti-tumor therapy.

20. Noteborn, M.H.M., and De Boer, G.F. (1996). Patent USA/no. 030, 335.

21. Noteborn, M.H.M., De Boer, G.F., Van Roozelaar, D., Karreman, C., Kranenburg, O., Vos, J., Jeurissen, S., Zantema, A., Hoebe, R., Koch, G., Van Ormondt, H., and Van der Eb, A.J. (1991). Characterization of cloned chicken anemia virus DNA that contains all elements for the infectious replication cycle. *Journal of Virology* 65, 3131-3139.

22. Noteborn, M.H.M., Hoebe, R.C., and Pietersen, A. (1997). A gene delivery vehicle expressing the apoptosis-inducing proteins VP2 and/or apoptin. European Patent Application no. 97201121.7

23. Noteborn, M.H.M., Todd, D., Verschueren, C.A.J., De Gauw, H.W.F.M., Curran, W.L., Veldkamp, S., Douglas, A.J., McNulty, M.S., Van der Eb, A.J., and Koch, G. (1994). A single chicken anemia virus protein induces apoptosis. *Journal of Virology* 68, 346-351.

24. Noteborn, M.H.M., and Zhang, Y. (1997). Methods and means for determining the transforming capability of agents, for determining the predisposition of cells to become transformed and prophylactic treatment of cancer using apoptin-like activity. European Patent Application no. 97439

25. Paulovich, A.G., Toczyski, D., Hartwell, H. (1997). When checkpoints fail. *Cell* 88, 315-321.

26. Rabbitts, T.H. (1991). *Cell* 67, 641-644.

27. Rose, M.D., Winston, F., and Hieter, P. (1990). *Methods in yeast genetics. A laboratory course manual*. Cold Spring Harbor Laboratory Press, Cold Spring Harbor, New York, USA.

28. Sachs, L. and Lotem, J. (1993). Control of programmed cell death in normal and leukemia cells: New implications for therapy. *Blood* 82, 15-21.

29. Sanger, F., Nicklen, S., and Coulson, A.R. (1977). DNA sequencing with chain-terminating inhibitors. *Proceedings National Academic Sciences USA* 74, 5463-5467.

30. Steller, H. (1995). Mechanisms and genes of cellular suicide. *Science* 267, 1445-1449.

31. Sawyers, C.L. and Denny, C.T. (1994) *Cell* 77, 171-173.

32. Telford, W.G., King, L.E., Fraker, P.J. (1992). Comparative evaluation of several DNA binding dyes in the detection of apoptosis-associated chromatin degradation by flow cytometry. *Cytometry* 13, 137-143.

33. Teodoro, J.G. and Branton, P.E. (1997). Regulation of apoptosis by viral gene products. *Journal of Virology* 71, 1739-1746.

34. Thompson, C.B. (1995). Apoptosis in the pathogenesis and treatment of disease. *Science* 267, 1456-1462.

35. White, E. (1996). Life, death, and the pursuit of apoptosis. *Genes and development* 10, 1-15.

36. Wyllie, A.H. (1995). The genetic regulation of apoptosis. *Current Opinion in Genetics and Development* 5, 97-104.

37. Wyllie, A.H., Kerr, J.F.R., Currie, A.R. (1980). Cell death: The significance of apoptosis. *International Review of Cytology* 68, 251-306.

38. Yang, X., Hubbard, E.J.A., and Carlson, M. (1992). A protein kinase substrate identified by the two-hybrid system. *Science* 257, 680-682.

39. Zhuang, S.-M., Landegent, J.E., Verschueren, C.A.J., Falkenburg, J.H.F., Van Ormondt, H., Van der Eb, A.J.,

## SEQUENCE LISTING

## (1) GENERAL INFORMATION:

## (i) APPLICANT:

- (A) NAME: Leadd B.V.
- (B) STREET: Wassenaarseweg 72
- (C) CITY: Leiden
- (D) STATE: Zuid-Holland
- (E) COUNTRY: the Netherlands
- (F) POSTAL CODE (ZIP): 2333 AL

(ii) TITLE OF INVENTION: Novel molecules involved in  
apoptotic pathways.

(iii) NUMBER OF SEQUENCES: 14

## (iv) COMPUTER READABLE FORM:

- (A) MEDIUM TYPE: Floppy disk
- (B) COMPUTER: IBM PC compatible
- (C) OPERATING SYSTEM: PC-DOS/MS-DOS
- (D) SOFTWARE: PatentIn Release #1.0, Version  
#1.30 (EPO)

## (v) CURRENT APPLICATION DATA:

APPLICATION NUMBER: EP 97203781.6

## (2) INFORMATION FOR SEQ ID NO: 1:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 17 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: unknown
- (D) TOPOLOGY: unknown

(ii) MOLECULE TYPE: other nucleic acid

(iii) HYPOTHETICAL: NO

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 1:

TACCACTACA ATGGATG  
17

## (2) INFORMATION FOR SEQ ID NO: 2:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 658 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: unknown
- (D) TOPOLOGY: unknown

(ii) MOLECULE TYPE: DNA (genomic)

## (ix) FEATURE:

- (A) NAME/KEY: misc feature  
 (B) LOCATION: 89..716  
 (D) OTHER INFORMATION: /label= N  
 /note= "N" stands for unknown."

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 3:

CGGAGTTACA AGAGGCTACC AAAGAATTCC AGATTAAAGA GGATATTCCT  
 GAAACAAAGA 60

TGAAATTCTT ATCAGTTGAA ACTCCTGANA ATGACAGCCA GTTGTCAAAT  
 ATCTCCTGTT 120

CGTTTCAAGG TGAGCTCGAA AGTTCCTTAT GAGATACAAA AAGGACAATG  
 CACTTATCAC 180

CTTTGAAAAA GGAAGAAGTT GCTCAAAATG TGNGTAANGC ATGAGTAAAC  
 ATCATGTACA 240

GATAATAAGA TGTAATCTG GAGGTTACGG CCAAAGCCAA GTTCCATTAA  
 TATTCAAGGA 300

GTCANGATTG CAGNGTTTAT GCTAGAANGT TTCTAAAAAT GANAATCAAT  
 GGTTACTGGA 360

AATTCCTGGA CACATTGCGN TGAAAGATCA AGATGACGAA GACAACTAA  
 GAAGCTGAGC 420

TTTTCAAAAG TCCCGAAANA TGGAAGAGCG GTAGAGGGTG GNACCGCGTG  
 NGANCTATGA 480

CAAGACAAGN CCGGGGAAGN TGCAGTCCAT CACGTTTGTN NGAAGATTGG  
 ANGTNGGCTG 540

ACCAANGAAT TTTGAAAAAG GAGANGAATT ACCCCTCTTT ANGAGTAANA  
 TCAAACCCT 600

GCCATAANAA GTTNACTGGT TTCNCCCATT ACACAGNANT TACANNTTGA  
 NCAANANTAN 660

NCAGGATAAT TTNCAGGGGA ANAATCTNAA GNATGGCAAG NTGACTTCTG  
 GACAANGGT 719

## (2) INFORMATION FOR SEQ ID NO: 4:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 220 amino acids  
 (B) TYPE: amino acid  
 (C) STRANDEDNESS: unknown  
 (D) TOPOLOGY: unknown

(ii) MOLECULE TYPE: protein

(iii) HYPOTHETICAL: NO

185

**5**

205

**10**

## 15

## 15

15

15

15

15

20

20

## 25

25

10

**30**

**30**

25

35

35

45

40

60

45

75

50

50

90

50

55



305

## (2) INFORMATION FOR SEQ ID NO: 6:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 659 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: unknown
- (D) TOPOLOGY: unknown

(ii) MOLECULE TYPE: DNA (genomic)

(iii) HYPOTHETICAL: NO

## (ix) FEATURE:

- (A) NAME/KEY: misc feature
- (B) LOCATION: 189..657
- (D) OTHER INFORMATION: /label= N  
/note= "N" stands for unknown."

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 6:

AGCAGGTGCT GCAACAAAG GAGCACACGA TCAACATGGA GGAGTGCCGG  
CTGCGGGTGC 60

AGGTCCAGCC CTTGGAGCTG CCCATGGTCA CCACCATCCA GGTGTCCAGC  
CAGTTGAGTG 120

GCCGGAGGGT GTTGGTCACT GGATTTCTTG CCAGCCTCAG GCTGAGTGAG  
GAGGAGCTGC 180

TGGACAANCT ANAGATCTTC TTTGGCAAGA CTAGGAACGG AGGTGGCNAT  
GTGGACNTTC 240

GGGANCTACT GCCAGGGANT GTCATGCTGG GGTTCGCTAG GGATGGAGTG  
GCTCANCCTC 300

TGTGCCAAAT CGGCCATTTC ACAGTGCCAC TGGGTGGGCA GCANGTCCCT  
CTGAGAGTCT 360

CTCCGTATGT GAATGGGGAN ATCCAGANGG CTGANATCAG GTCNCAGCCA  
NTTCCCCGCT 420

CGGTACTGGT GCTCAACATT CCTGATATCT TGGATGGCCC GGAGCTGCAT  
GACGTCCTGG 480

ANATCCACTT CCAGAANCCC ACCCGCGGGG GCGGAGATGT AAGACGCCCT  
GACAGTCGTA 540

CCCCAAGGAC AACAGGGCCT AACAGTCTTC ACCTCCTGAA TCAAGGCTAN  
GGGCCTCCCC 600

CTTCTCATCC TCCCCACCCC CCCC GCCAAA GGTTCTCAAN ACTGGGCCTG  
GGCTTTNTG 659

## (2) INFORMATION FOR SEQ ID NO: 7:

- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: unknown
- (D) TOPOLOGY: unknown

(ii) MOLECULE TYPE: DNA (genomic)

(iii) HYPOTHETICAL: NO

(ix) FEATURE:

- (A) NAME/KEY: misc\_feature
- (B) LOCATION: 182..626
- (D) OTHER INFORMATION: /label= N  
/note= "N" stands for unknown."

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 8:

GGATCCACTG CCCTCTGCTT GCGGGCTCTG CTCTGATCAC CTTTGATGAC  
CCCAAAGTGG 60

CTGAGCAGGT GCTGCAACAA AAGGAGCACA CGATCAACAT GGAGGAGTGC  
CGGCTGCGGG 120

TGCAGGTCCA GCCCTTGGAG CTGCCCATGG TCACCACCAT CCAGGTGATG  
GTGTCCAGCC 180

ANITGAGTGG CCGGAGGGTG TTGGTCACTG GATTTCCTGC CAGCCTCAGG  
CTGANTGAGG 240

AGGAGCTGCT GGACAAGCTA TGAGATCTTC TTTGGCAANA CTANGAACGG  
ANGTGGCGAT 300

GTGGACGTTT GGGAGCTACT GCCAGGGAGT GTCATGCTGG GGTTTGCTAC  
GGATGGAGTG 360

GCTCAGCGTC TGTGCCAAAT CGGCCAGTTC ACAAGTGCCA CTGGGTGGGC  
AGCAAGTCCC 420

TCTGAGAGTC TCTCCGTATG TGANTGGNGA GATCAGAATG CTGANATTAA  
GTCGCATCCA 480

ATTCTCTGCT CNGGTACTGG TGCTCANNAT CCTGANATCT TGGATTGGCC  
CCNGANTNCA 540

TGANATCTGG NAGATTCAAT TNCANAAGTC CANCCNNCNG NGNCGGGAAG  
TANANGCCCG 600

ANANTTCNTN NCNTANGGNC AGCANNGCCT G  
631

(2) INFORMATION FOR SEQ ID NO: 9:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 138 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS: unknown
- (D) TOPOLOGY: unknown

(ii) MOLECULE TYPE: protein

(iii) HYPOTHETICAL: NO

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 10:

Met Ser Ala Pro Leu Asp Ala Ala Leu His Ala Leu Gln  
 Glu Glu Gln  
 1 5 10  
 15

Ala Arg Leu Lys Met Arg Leu Trp Asp Leu Gln Gln Leu  
 Arg Lys Glu  
 20 25  
 30

Leu Gly Asp Ser Pro Lys Asp Lys Val Pro Phe Ser Val  
 Pro Lys Ile  
 35 40 45

Pro Leu Val Phe Arg Gly His Thr Gln Gln Asp Pro Glu  
 Val Pro Lys  
 50 55 60

Ser Leu Val Ser Asn Leu Arg Ile His Cys Pro Leu Leu  
 Ala Gly Ser  
 65 70 75  
 80

Ala Leu Ile Thr Phe Asp Asp Pro Lys Val Ala Glu Gln  
 Val Leu Gln  
 85 90  
 95

Gln Lys Glu His Thr Ile Asn Met Glu Glu Cys Arg Leu  
 Arg Val Gln  
 100 105  
 110

Val Gln Pro Leu Glu Leu Pro Met Val Thr Thr Ile Gln  
 Val Ser Ser  
 115 120 125

Gln Leu Ser Gly Arg Arg Val Leu Val Thr Gly Phe Pro  
 Ala Ser Leu  
 130 135 140

Arg Leu Ser Glu Glu Glu Leu Leu Asp Lys Leu Glu Ile  
 Phe Phe Gly  
 145 150 155  
 160

Lys Thr Arg Asn Gly Gly Gly Asp Val Asp Val Arg Glu  
 Leu Leu Pro

## 50

- 55

Asp Leu Ser Leu Lys Ile Pro Glu Ile Ser Ile Gln Asp  
Met Thr Ala

165

170

175

Gln Val Thr Ser Pro Ser Gly Lys Thr His Glu Ala Glu  
Ile Val Glu

180

185

190

Gly Glu Asn His Thr Tyr Cys Ile Arg Phe Val Pro Ala  
Glu Met Gly

195

200

205

Thr His Thr Val Ser Val Lys Tyr Lys Gly Gln His Val  
Pro Gly Ser

210

215

220

Pro Phe Gln Phe Thr Val Gly Pro Leu Gly Glu Gly Gly  
Ala His Lys

225

230

235

240

Val Arg Ala Gly Gly Pro Gly Leu Glu Arg Ala Glu Ala  
Gly Val Pro

245

250

255

Ala Glu Phe Ser Ile Trp Thr Arg Glu Ala Gly Ala Gly  
Gly Leu Ala

260

265

270

Ile Ala Val Glu Gly Pro Ser Lys Ala Glu Ile Ser Phe  
Glu Asp Arg

275

280

285

Lys Asp Gly Ser Cys Gly Val Ala Tyr Val Val Gln Glu  
Pro Gly Asp

290

295

300

Tyr Glu Val Ser Val Lys Phe Asn Glu Glu His Ile Pro  
Asp Ser Pro

305

310

315

320

Phe Val Val Pro Val Ala Ser Pro Ser Gly Asp Ala Arg  
Arg Leu Thr

325

330

335

Val Ser Ser Leu Gln Glu Ser Gly Leu Lys Val Asn Gln  
Pro Ala Ser

340

345

350

Phe Ala Val Ser Leu Asn Gly Ala Lys Gly Ala Ile Asp  
Ala Lys Val

Val Ala Lys Gly Leu Gly Leu Ser Lys Ala Tyr Val Gly  
Gln Lys Ser

565

570

575

Ser Phe Thr Val Asp Cys Ser Lys Ala Gly Asn Asn Met  
Leu Leu Val

580

585

590

Gly Val His Gly Pro Arg Thr Pro Cys Glu Glu Ile Leu  
Val Lys His

595

600

605

Val Gly Ser Arg Leu Tyr Ser Val Ser Tyr Leu Leu Lys  
Asp Lys Gly

610

615

620

Glu Tyr Thr Leu Val Val Lys Trp Gly His Glu His Ile  
Pro Gly Ser

625

630

635

640

Pro Tyr Arg Val Val Val Pro  
645

(2) INFORMATION FOR SEQ ID NO: 13:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 213 amino acids

(B) TYPE: amino acid

(C) STRANDEDNESS: unknown

(D) TOPOLOGY: unknown

(ii) MOLECULE TYPE: protein

(iii) HYPOTHETICAL: NO

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 13:

His Glu Gly Arg Gly Val Thr Gly Asn Pro Ala Glu Phe  
Val Val Asn

1

5

10

15

Thr Ser Asn Ala Gly Ala Gly Ala Leu Ser Val Thr Ile  
Asp Gly Pro

20

25

30

Ser Lys Val Lys Met Asp Cys Gln Glu Cys Pro Glu Gly  
Tyr Arg Val

35

40

45

(iii) HYPOTHETICAL: NO

5

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 14:

10

His Glu Gly Arg Pro Thr Glu Pro Gly Asn Tyr Ile Ile  
 Asn Ile Lys  
 1 5 10  
 15

15

Phe Ala Asp Gln His Val Pro Gly Ser Pro Phe Ser Val  
 Lys Val Thr  
 20 25  
 30

20

Gly Glu Gly Arg Val Lys Glu Ser Ile Thr Arg Arg Arg  
 Arg Ala Pro  
 35 40 45

25

Ser Val Ala Asn Val Gly Ser His Cys Asp Leu Ser Leu  
 Lys Ile Pro  
 50 55 60

30

Glu Ile Ser Ile Gln Asp Met Thr Ala Gln Val Thr Ser  
 Pro Ser Gly  
 65 70 75  
 80

Lys Thr His Glu Ala Glu Ile Val Glu Gly Glu Asn His  
 Thr Tyr Cys  
 85 90  
 95

35

Ile Arg Phe Val Pro Ala Glu Met Gly Thr His Thr Val  
 Ser Val Lys  
 100 105  
 110

40

Tyr Lys Gly Gln His Val Pro Gly Ser Pro Phe Gln Phe  
 Thr Val Gly  
 115 120 125

Pro Leu Gly Glu Gly Gly Ala His Xaa Val Arg Ala Gly  
 Gly Pro Gly  
 130 135 140

45

Leu Xaa Lys Ser Ser Trp Ser Ala Ser Arg Ile Gln Tyr  
 Leu Gly Pro  
 145 150 155  
 160

50

Gly Lys Leu Val Leu Glu Ala Trp Pro Leu Leu Ser Xaa  
 Ala Pro Ala  
 165 170  
 175

55

induction of apoptosis in a population of cells related to a pathological condition.

14. An Nmi/Hou-like proteinaceous substance having at least a functional and/or specific part of the sequence of figure 3 or being encoded by a functional and/or specific part of the sequence of figure 1 or figure 2 or being at least 60, preferably 70, preferably 80% homologous to at least a functional and/or specific part of the sequence of figure 3 or being at least 60, preferably 70, preferably 80% homologous to a protein encoded by at least a functional and/or specific part of the sequence of figure 1 or figure 2.
15. A recombinant or isolated proteinaceous substance comprising at least a functional part of a member of the family of Nmi/Hou-like proteins or at least a functional part of a member of the family of Hou-like proteins for use in the induction of apoptosis in a population of cells related to a pathological condition.
16. An IFP35-like proteinaceous substance having at least a functional and/or specific part of the sequence of figure 6 or 7 or being encoded by a functional and/or specific part of the sequence of figure 4 or figure 5 or being at least 60, preferably 70, preferably 80% homologous to at least a functional and/or specific part of the sequence of figure 6 or 7 or being at least 60, preferably 70, preferably 80% homologous to a protein encoded by at least a functional and/or specific part of the sequence of figure 4 or figure 5.
17. A method for inducing apoptosis in cells comprising providing said cells with Nmi/Hou-like protein activity and/or IFP-35-like activity together with apoptin-like activity.
18. Use of apoptin to find proteinaceous substances associated with apoptosis.



CGGAGTTACAAGAGGCTACCAAAGAATTCCAGATTAAAGAGGATATTCCTGAAACAAAGATGAAA  
TTCTTATCAGTTGAAACTCCTGANAATGACAGCCAGTTGTCAAATATCTCCTGTTCGTTTCAAGG  
TGAGCTCGAAAGTTCCTTATGAGATACAAAAGGACAATGCACTTATCACCTTTGAAAAGGAAG  
AAGTTGCTCAAATGTGNGTAANGCATGAGTAAACATCATGTACAGATAATAAGATGTAAATCTG  
GAGGTTACGGCCAAAGCCAAGTTCATTAATATTCAAGGAGTCANGATTCCAGNGTTTATGCTAG  
AANGTTTCTAAAAATGANAATCAATGGTTACTGGAAATTCCTGGACACATTGCGNTGAAAGATCA  
AGATGACGAAGACAACTAAGAAGCTGAGCTTTTCAAAGTCCCGAAANATGGAAGAGCGGTAGA  
GGGTGGNACCGCGTGNGANCTATGACAAGACAAGNCCGGGGAAGNTGCAGTCCATCACGTTTGTN  
NGAAGATTGGANGTNGGCTGACCAANGAATTTTGAAAAGGAGANGAATTACCCCTCTTTANGAG  
TAANATCAAACCTGCCATAANAAGTTNACTGGTTTCNCCCATTACACAGNAN  
TTACANNTGANCAANANTANNCAGGATAATTTNCAGGGGAANAATCTNAAGNATGGCAAGNTGA  
CTTCTGGACAANGGT

**Figure 2**

Hou c17/#2

AGCAGGTGCTGCAACAAAAGGAGCACACGATCAACATGGAGGAGTGCCGGCTGCGGGTGCAGGTC  
CAGCCCTTGAGCTGCCCATGGTCACCACCATCCAGGTGTCCAGCCAGTTGAGTGGCCGGAGGGT  
GTTGGTCACTGGATTTCCTGCCAGCCTCAGGCTGAGTGAGGAGGAGCTGCTGGACAANCTANAGA  
TCTTCTTTGGCAAGACTAGGAACGGAGGTGGCNATGTGGACNTTCGGGANCTACTGCCAGGGANT  
GTCATGCTGGGGTTTGCTAGGGATGGAGTGGCTCANCCTGTGTGCCAAATCGGCCATTTACAGT  
GCCACTGGGTGGGCAGCANGTCCCTCTGAGAGTCTCTCCGTATGTGAATGGGGANATCCAGANGG  
CTGANATCAGGTCNCAGCCANTTCCCCGCTCGGTACTGGTGCTCAACATTCCTGATATCTTGGAT  
GGCCCGGAGCTGCATGACGTCCTGGANATCCACTCCAGAANCCCACCCGCGGGGGCGGAGATGT  
AAGACGCCCTGACAGTCGTACCCCAAGGACAACAGGGCCTAACAGTCTTCACCTCCTGAATCAAG  
GCTANGGGCCTCCCCCTTCTCATCCTCCCCACCCCCCGCCAAAGGTTCTCAANACTGGGCCTG  
GGCTTTNTG

**Figure 4**

IFP35 c14/#1

GGATCCACTGCCCTCTGCTTGCGGGCTCTGCTCTGATCACCTTTGATGACCCCAAAGTGGCTGAG  
CAGGTGCTGCAACAAAAGGAGCACACGATCAACATGGAGGAGTGCCGGCTGCGGGTGCAGGTCCA  
GCCCTTGGAGCTGCCCATGGTCACCACCATCCAGGTGATGGTGTCCAGCCANTTGAGTGGCCGGA  
GGGTGTTGGTCACTGGATTTCTGCCAGCCTCAGGCTGANTGAGGAGGAGCTGCTGGACAAGCTA  
TGAGATCTTCTTTGGCAANACTANGAACGGANGTGGCGATGTGGACGTTTCGGGAGCTACTGCCAG  
GGAGTGTCATGCTGGGGTTTGTACGGATGGAGTGGCTCAGCGTCTGTGCCAAATCGGCCAGTTC  
ACAAGTGCCACTGGGTGGGCAGCAAGTCCCTCTGAGAGTCTCTCCGTATGTGANTGGNGAGATCA  
GAATGCTGANATTAAGTCGCATCCAATTCCTCGCTCNGGTACTGGTGCTCANNATCCTGANATCT  
TGGATTGGCCCCNGANTNCATGANATCTGGNAGATTCAATTNCANAAGTCCANCCNNCNGNGNCG  
GGAAGTANANGCCCGANANTTCNTNNCNTANGGNCAGCANNGCCTG

**Figure 6**

IFP35 c51/#3

Filamin	1	RLRNQHVGISFVPKETGEHLVHVKNQGQHVASSPIPVVISQSEIGDASRVVSGQGLHEG
c50/01	1	-----
c57/02	1	-----
Filamin	61	HTFEPAEFIIDTRDAGYGGLSLSLIEGFSKVDINTEDLEDGTCRVTYCPTTEPGNYIINIKF
c50/01	1	-----
c57/02	1	-----HEGRPTTEPGNYIINIKF
Filamin	121	ADQHVPGSPFSVKVTGEGRVKESITRRRRAPSVANVGSHCDLSLKIPEISIQDMTAQVTS
c50/01	1	-----
c57/02	18	ADQHVPGSPFSVKVTGEGRVKESITRRRRAPSVANVGSHCDLSLKIPEISIQDMTAQVTS
Filamin	181	PSGKTHEAEIVEGENHTYTCIRFVPAEMGTHTVSVKYKGOHVPGSPFQFTVGPLGEGGAHX
c50/01	1	-----
c57/02	78	PSGKTHEAEIVEGENHTYTCIRFVPAEMGTHTVSVKYKGOHVPGSPFQFTVGPLGEGGAHX
Filamin	241	VRAGGPGLEEEFGVPFEFS.EWTREAGAGLAJAVEPEKAEISFEDRDSCGAYV
c50/01	1	-----
c57/02	138	VRAGGPGLEEEFGVSAEIRIQYCGPKLVLEWFLSXPEKLISLLRTATPVVLMV
Filamin	300	QEPGDYEVSVKFNEEHIPDSFFVVPVASPSGDARRLTVSSSLQESOLKVNQPASFAVSLNG
c50/01	1	-----
c57/02	197	XEPSD*INPXQVSTKEHX-----
Filamin	360	AKGAIDAKVHSPSGALEECYVTEIDQDKYAVRFIPRENGVYLIDVKFNGTHIPGSPFKIR
c50/01	1	-----
c57/02	214	-----
Filamin	420	VGEPPGHGGDPGLVSAYGAGLEG.GVTGNPAEFVVNTSNAGAGALSVTIDGPSKVKHDCQE
c50/01	1	-----HEGRGVTGNPAEFVVNTSNAGAGALSVTIDGPSKVKHDCQE
c57/02	214	-----
Filamin	479	CPEGYRVITYTPMAPGSYLISIKYGGPYHIGGSPFKAKVTGPRLVSNHSLHETSSVFVDSL
c50/01	42	CPEGYRVITYTPMAPGSYLISIKYGGPYHIGGSPFKAKVTGPRLVSNHSLHETSSVFVDSL
c57/02	214	-----
Filamin	539	TKATCAPQHGAPGPGPADASKVVAKGLGLSKAYVQOKSSFTVDCSKAONNMLLVGVHGP
c50/01	102	TKATCAPQHGAPGPGPADASKVVAKGLGLSKAYVCEKSSFTVDCSKACIIMLLVGVHGP
c57/02	214	-----
Filamin	599	TPCEILVKEVGS.RDYSVSYLLKDKGE.YTLVVKWGHEHIEGSEYRVVP-
c50/01	162	TPCEILVKARGQPAQORVLTCPKDKGEVHTGGQNGCDYQLECKELPCCGP
c57/02	214	-----

Figure 8



European Patent  
Office

## PARTIAL EUROPEAN SEARCH REPORT

Application Number  
EP 97 20 3781

DOCUMENTS CONSIDERED TO BE RELEVANT			CLASSIFICATION OF THE APPLICATION (Int.CL.6)
Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	
A	ZHUANG S -M ET AL: "APOPTIN, A PROTEIN ENCODED BY CHICKEN ANEMIA VIRUS, INDUCES CELL DEATH IN VARIOUS HUMAN HEMATOLOGIC MALIGNANT CELLS IN VITRO" LEUKEMIA, vol. 9, no. SUPPL. 01, October 1995, pages S118-S120, XP000602147 * the whole document *	1-7,9-15	
A	ZHUANG S -M ET AL: "APOPTIN, A PROTEIN DERIVED FROM CHICKEN ANEMIA VIRUS, INDUCES P53- INDEPENDENT APOPTOSIS IN HUMAN OSTEOSARCOMA CELLS" CANCER RESEARCH, vol. 55, no. 3, 1 February 1995, pages 486-489, XP000602162 * the whole document *	1-7,9-15	
T	DE 196 28 894 A (HAGENMAIER HANS PAUL) 22 January 1998 * claims 1-14,16,17 *	1,13	TECHNICAL FIELDS SEARCHED (Int.CL.6)

EPO FORM 1503 03/93 (P04C10)

**This Page is Inserted by IFW Indexing and Scanning  
Operations and is not part of the Official Record**

**BEST AVAILABLE IMAGES**

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

- ☐ BLACK BORDERS
- ☒ IMAGE CUT OFF AT TOP, BOTTOM OR SIDES
- ☐ FADED TEXT OR DRAWING
- ☐ BLURRED OR ILLEGIBLE TEXT OR DRAWING
- ☐ SKEWED/SLANTED IMAGES
- ☐ COLOR OR BLACK AND WHITE PHOTOGRAPHS
- ☐ GRAY SCALE DOCUMENTS
- ☒ LINES OR MARKS ON ORIGINAL DOCUMENT
- ☐ REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY
- ☐ OTHER: \_\_\_\_\_

**IMAGES ARE BEST AVAILABLE COPY.**

**As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.**